

AD-A153 665

LASER-STIMULATED VIBRATIONAL EXCITATION OF AN ADSPECIES 1/1  
STUDIED BY A GENE. (U) ROCHESTER UNIV NY DEPT OF  
CHEMISTRY A C BERI ET AL APR 85

UNCLASSIFIED

UROCHESTER/DC/85/TR-62 N00014-80-C-0472


F/G 20/5

NL

END

FILED

ONE



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

(2)

AD-A153 665

OFFICE OF NAVAL RESEARCH  
Contract N00014-80-C-0472  
Task No. NR 056-749  
TECHNICAL REPORT No. 62

Laser-Stimulated Vibrational Excitation of an Adsorbed Species  
Studied by a Generalized Master Equation: Neutral  
Atomic Hydrogen on Hydrated KCl(001)

by

A. C. Beri and Thomas F. George

Prepared for Publication

in

Journal of Vacuum Science and Technology B

Department of Chemistry  
University of Rochester  
Rochester, New York 14627

April 1985

DTIC  
ELECTE  
MAY 13 1985  
B

DTIC FILE COPY

Reproduction in whole or in part is permitted for  
any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

85 04 15 008

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UROCHESTER/DC/85/TR-62			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry University of Rochester		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research (Code 413)		
6c. ADDRESS (City, State and ZIP Code) River Station Rochester, New York 14627		7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-80-C-0472		
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO. 61153N	PROJECT NO. 013-08	TASK NO. NR 056-749	WORK UNIT NO.
11. TITLE Laser-Stimulated Vibrational Excitation of an Adsorbed Species Studied by a Generalized Master Equation: Neutral Atomic Hydrogen on Hydrated KCl(001)					
12. PERSONAL AUTHOR(S) A. C. Beri and Thomas F. George					
13a. TYPE OF REPORT Interim Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) April 1985	
15. PAGE COUNT 21					
16. SUPPLEMENTARY NOTATION Prepared for publication in Journal of Vacuum Science and Technology B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	VIBRATIONAL EXCITATION, GENERALIZED MASTER EQUATION, LASER-STIMULATED, QUANTUM TREATMENT, ADSORBED SPECIES, H(H <sub>2</sub> O)/KCl(001) SYSTEM		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Vibrational energy transfer between an IR laser, a solid surface and an adatom is studied by a generalized master equation which includes memory effects. Numerical problems associated with the temporally delocalized memory kernel are overcome by introducing the isomnesic approximation. Results obtained for time scales ranging from 2 ps to 2 μs for H(H <sub>2</sub> O)/KCl(001) show that effective vibrational excitation is possible with low power lasers (1 W/cm <sup>2</sup> ) in spite of fast phonon relaxation. Detailed time-evolution of the system is seen to differ radically from that predicted by Markovian theories, even for fairly long time scales.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

## Abstract

Vibrational energy transfer between an IR laser, a solid surface and an adatom is studied by a generalized master equation which includes memory effects. Numerical problems associated with the temporally delocalized memory kernel are overcome by introducing the isomnesic approximation. Results obtained for time scales ranging from 2 ps to 2  $\mu$ s for H(H<sub>2</sub>O)/KCl(001) show that effective vibrational excitation is possible with low power lasers (1 W/cm<sup>2</sup>) in spite of fast phonon relaxation. Detailed time-evolution of the system is seen to differ radically from that predicted by Markovian theories, even for fairly long time scales.

A session for  
NIGHT CRA&I ☒  
DAY BAR ☐  
MORNING BAR ☐  
CITY BAR ☐

A-1

## I. Introduction

Developments in the experimental study of clean, well-characterized solid surfaces and surface processes such as scattering, adsorption, desorption and migration in the presence of laser radiation have resulted in important technological applications (e.g., laser-assisted vapor deposition of microcircuits). The underlying dynamics of these heterogeneous systems is a many-body problem involving the elementary excitations of the solid and its surface, such as phonons, electrons and plasmons, the degrees of freedom of the adsorbed species or gas molecules, and those of the laser.<sup>1,2</sup> The excitation process can be considered as direct laser-pumping of the substrate leading to a primarily non-selective thermal effect, or a pumping of the adsorbed species, which can be selective in the sense that only a single degree of freedom is excited by the laser. The resulting energy absorption and the dissipation involving all other degrees of freedom govern the overall chemical process.<sup>3-8</sup>

In this work, the vibrational excitation of a prototype system, viz. the adsorptive bond (adbond) between an atom and a surface is studied. This system is not in the same category as those in which internal vibrational modes of an adsorbed molecule are excited, with subsequent energy transfer to the adbond.<sup>7,9</sup> It is substantially simpler, in view of the aim of this work, which is to investigate the mechanisms involved in IR laser-stimulated surface processes (LSSP) using a treatment which is as close to first-principles as possible. One of the fundamental questions addressed is whether a low-power laser, because of its coherence, can deposit large amounts of energy into a specific mode of a system (such as an adbond) which is coupled to energy sinks, such as phonons, which are not infinite.

Early classical treatments<sup>10</sup> of the problem and phenomenological quantum treatments<sup>4,5</sup> provided conflicting answers to this question. Recent, more fundamental studies<sup>8</sup> have suggested that extremely high laser powers would be needed to effectively desorb an adspecies. However, our preliminary work<sup>11</sup> showed that a very small degree of detuning between the laser and energy levels of the system excluded the possibility of long-term energy absorption. The reason for this behaviour is seen to be the oscillatory nature of the laser energy absorption "rate" (actually, the memory kernel) for the adbond, with a frequency related to the detuning, typically a small percentage of the laser frequency itself, viz.  $\sim 10^{10}$ - $10^{11}$  s<sup>-1</sup>. Thus, for processes lasting over a few hundred picoseconds, substantial cancellations in the overall absorption can be expected, and only a very high-power laser would be able to compete with the loss of energy from the adbond to phonons through relaxation effects. In view of this, the systems chosen here have little or no detuning, viz. systems with one bound-to-bound transition exactly resonant with the laser frequency and another very close to or equal to the Debye frequency of the solid. This represents an ideal situation which emphasizes the coherent nature of laser radiation.

The widely varying temporal behaviour of dynamical systems far from equilibrium is well established. From a fundamental point of view, theoretical treatments of such systems depend crucially on the time scales being considered. The most common long-time limit, for example, allows a description of energy transfer in terms of average relaxation rates in the Markovian approximation, where approach to equilibrium is assumed to be monotonic. Non-Markovian behaviour is assumed to last for extremely short times compared to typical relaxation times of the system. The forces are considered to be random, with a very short correlation time. Resonant excitation with a low-power laser does not satisfy these conditions, and

one should expect non-Markovian behaviour to persist for times that increase with decrease in laser power. Whether the two approaches lead to the same steady state is also open to question.

Non-Markovian effects are conveniently treated by using a generalized master equation (GME) to describe the time-evolution of the adbond (the "relevant" system) with the laser and phonon fields as two uncoupled irrelevant systems (reservoirs).<sup>12-14</sup> The Liouville equation for such a system is conveniently solved by using the Zwanzig projection-operator technique<sup>12</sup> and results in a set of coupled integrodifferential equations for the occupation probabilities  $P_S$  of states  $|S\rangle$  of the adbond:<sup>14</sup>

$$\dot{P}_S(t) = \sum_{S' \neq S} \int_0^t dt' [K_{SS'}(t-t')P_{S'}(t') - K_{S'S}(t-t')P_S(t')] , \quad (1)$$

where the memory kernels  $K_{SS'}$  include contributions due to the phonon field  $K^{(p)}$  and the radiation field  $K^{(r)}$ :

$$K_{SS'}(t) = K_{SS'}^{(p)}(t) + K_{SS'}^{(r)}(t). \quad (2)$$

Using a one-dimensional lattice and a Debye model to describe the lattice vibrations, we have obtained  $K_{SS'}^{(p)}$  in a closed form which involves a summation over all lattice positions  $\ell$  and number of phonons  $n$  independently.<sup>11,15</sup> The former is due to a summation of pair potentials between the adatom and the lattice atoms, while the latter results from an expansion of  $\exp[y_{\ell\ell'}(t)]$  in a Taylor series, where  $y_{\ell\ell'}(t)$  is the lattice displacement correlation function

$$y_{\ell\ell'}(t) = \langle\langle u_{\ell}(t)u_{\ell'} \rangle\rangle . \quad (3)$$

Using techniques described elsewhere,<sup>11</sup>  $y_{\ell\ell'}(t)$  can be obtained in closed form as

$$y_{\ell\ell'}(t) = R_{\ell\ell'}(t) + i I_{\ell\ell'}(t) , \quad (4)$$



where  $R_{\ell\ell'}$  and  $I_{\ell\ell'}$  involve products of polynomials in  $t$  and  $\cos(\omega_D t)$  or  $\sin(\omega_D t)$ , and  $\omega_D$  is the Debye frequency. The phonon part of the memory kernel can then be written in terms of all the system parameters as follows:

$$\begin{aligned} K_{SS'}^{(p)}(t) = & A_p \sum_n (B_p)^n \sum_{\ell\ell'} \sum_{\mu\mu'} (\mu\mu')^n \\ & \times \lambda_{\ell\ell'\mu\mu'}(S+S') [\tilde{E}_{n\ell\ell'}(t) \cos(\omega_{SS'} t) + \tilde{O}_{n\ell\ell'}(t) \sin(\omega_{SS'} t)]. \end{aligned} \quad (5)$$

Here

$$A_p = 8 \left( \frac{D_e^{(0)}}{\hbar} \right)^2, \quad (6)$$

$$B_p = 6 (\beta^{(0)} k_B T)^2 / m_S \hbar \omega_D^3, \quad (7)$$

$$\tilde{E}_{n\ell\ell'}(t) = \sum_{j_e=0,2,4,\dots} \frac{(-1)^{j_e/2}}{j_e! (n-j_e)!} R_{\ell\ell'}^{n-j_e}(t) I_{\ell\ell'}^{j_e}(t), \quad (8)$$

$$\tilde{O}_{n\ell\ell'}(t) = \sum_{j_o=1,3,5,\dots} \frac{(-1)^{(j_o-1)/2}}{j_o! (n-j_o)!} R_{\ell\ell'}^{n-j_o}(t) I_{\ell\ell'}^{j_o}(t), \quad (9)$$

$$\max(j_e, j_o) \leq n. \quad (10)$$

$B_p$  is a smallness parameter which dictates, in part, the convergence properties of the expansion in number of phonons  $n$ . It is of the order of .02 for common temperatures  $T$  and common Debye frequencies ( $k_B$  is the Boltzmann constant and  $m_S$  the mass of a lattice atom).  $\mu$  and  $\mu'$  take on the values 1 and 2. The supermatrix  $\lambda$  involves powers of  $\exp(a)$  and  $\exp(\langle u^2 \rangle)$  and the matrix elements

$$E_{SS'}^{(\mu)} = \langle S | e^{\mu(z-z_0^{(0)})} | S' \rangle \quad (11)$$

where  $a$  is the lattice constant,  $u$  is the lattice displacement operator, and  $z$  is the distance of the adatom from the equilibrium position of the outermost lattice atom. The frequencies  $\omega_{SS'}$  correspond to transitions between levels  $|S\rangle$  and  $|S'\rangle$  of the adbond represented by a Morse potential with depth  $D_e^{(0)}$ , equilibrium distance  $z_0^{(0)}$  and exponent  $\beta^{(0)}$ .

The radiation part  $K_{SS'}^{(r)}$  of the memory kernel is given by

$$K_{SS'}^{(r)}(t) = A_r z_{S'S} z_{SS'} \cos(\omega_L - |\omega_{SS'}|)t, \quad (12)$$

where  $z_{SS'} = \langle S|z|S'\rangle$ . The amplitude  $A_r = (Ie_0^2/\epsilon_0\hbar^2)$  is proportional to the laser frequency  $I$  and the square of the charge separation  $e_0$  between the adspecies and the surface.

The complicated form of  $R_{\ell\ell'}$  and  $I_{\ell\ell'}$  precludes analytic solution of the GME. Numerical solution is made difficult by the presence of high frequencies in  $K_{SS'}^{(p)}$  and the large difference in the magnitudes of  $A_r$  ( $\sim 10^{17} \text{ s}^{-2}$  for a  $1 \text{ W/cm}^2$  laser) and  $A_p$  ( $\sim 10^{29} \text{ s}^{-2}$  for  $D_e^{(0)} = 0.1 \text{ eV}$ ). One can only solve the exact GME for a few periods of oscillation of  $K_{SS'}$ , which amounts to a few picoseconds.

Examination of a typical phonon kernel, Fig. 1, shows that it has substantial amplitude only for times of the order of a Debye period  $\tau_D$ , representative of phonon correlation times. The laser kernel, on the other hand, has constant amplitude for the case  $\omega_L = \omega_{SS'}$  for a given pair of levels  $(S, S')$ . For times larger than  $\tau_D$ , the integrated effect due to  $K_{SS'}^{(p)}$  will be unchanging because of cancellations, whereas that due to  $K_{SS'}^{(r)}$  will be cumulative. We have recently demonstrated this effect<sup>11</sup> using the exact GME (with a very high laser power in order for the effect to manifest itself in the very short time for which solution was possible), and predicted, via extrapolation arguments, that effective pumping of the

adbond with a low-power laser would be possible if the long-time regime could be examined. We present here an actual calculation of this long-time behaviour using low-power lasers, and find that the average adbond energy, given by

$$E(t) = \sum_S E_S P_S(t) \quad (13)$$

where  $E_S$  are eigenvalues of the zeroth order adbond Hamiltonian, rises rapidly after an initial loss to the phonon field, and exhibits a decidedly non-Markovian pattern throughout.

The approximations required for the long-time treatment are described in Section II. The resulting probability and energy profiles are presented in Section III, where comparison is made with the exact results (for a short-time scale) and with a Markovian limit.

## II. The Isomnesic (Constant-Memory) Approximation

We begin with the assertion that the physically important situation is one where the laser frequency and the Debye frequency equal different transition frequencies  $\omega_{SS'}$  of the adbond.<sup>11</sup> For this case  $K_{SS'}^{(r)}(t)$  becomes a constant  $k_{SS'}$  (from the term with  $|\omega_{SS'}| = \omega_L$ ) plus oscillating terms which we ignore in the sense of the rotating-wave approximation.<sup>16</sup> The phonon term,  $K_{SS'}^{(p)}(t)$ , has a complicated behaviour typified by  $K_{03}^{(p)}(t)$  in Fig. 1, but we note that it has appreciable magnitude for times  $\leq 2$  ps, and almost vanishes subsequently. If the mesh size of our theoretical experiment is no less than 2 ps, we can treat  $K_{SS'}^{(p)}(t)$  as a delta function, so that, from Eq. (2),

$$K_{SS'}(t) = \Omega_{SS'} \delta(t) + k_{SS'} \quad (14)$$

where

$$\Omega_{SS'} = \int_0^\infty dt K_{SS'}^{(p)}(t) \quad (15)$$

In Eq. (14), the first term, originating from the coupling of the adbond to the phonons, is the amnesic or Markovian term, and would lead, in the absence of the laser term, to the elimination of all memory effects. The second term, due to the resonant laser coupling, is the isomnesic non-Markovian term, and represents a constant memory effect all the way back to  $t=0$ .

The subsequent treatment is simplified by defining "diagonal" elements  $\Omega_{SS}$  as

$$\Omega_{SS} = - \sum_{S' \neq S} \Omega_{SS'} \quad (16)$$

and similarly for  $k_{SS}$ , and by introducing the vector notation  $\underline{P}(t) \equiv \{P_S(t)\}$  and matrix notation  $\underline{\Omega} \equiv \{\Omega_{SS'}\}$ ,  $\underline{k} \equiv \{k_{SS'}\}$ , etc. The exact GME may be written as

$$\dot{\underline{P}}(t) = \int_0^t dt' \underline{k}(t-t') \cdot \underline{P}(t') ; \quad (17)$$

in the isomnesic approximation (IA) it becomes

$$\dot{\underline{P}}(t) = \underline{k} \cdot \int_0^t dt' \underline{P}(t') + \underline{\Omega} \cdot \underline{P}(t) . \quad (18)$$

Formally this can be solved without any further approximations. Differentiating both sides of Eq. (18) we get the second-order differential equation

$$\ddot{\underline{P}}(t) = \underline{k} \cdot \underline{P}(t) + \underline{\Omega} \cdot \underline{P}(t) , \quad (19)$$

whose Laplace transform,

$$\underline{M}(s) \cdot \underline{P}(s) = \underline{I}(s) , \quad (20)$$

where

$$\underline{M}(s) \equiv \underline{k} + s\underline{\Omega} - s^2 \underline{\Delta} , \quad (21)$$

$$\underline{I}(s) \equiv -s\underline{P}(0) . \quad (22)$$

and  $\underline{P}(s)$  is the Laplace transform of  $\underline{P}(t)$ , is easily inverted using the Heaviside expansion theorem.

While this effectively eliminates the non-Markovian bottleneck, it is instructive to generate solutions of a Markovian version of this class of

problems. To do this, we have to make provisions for localizing  $K_{SS'}^{(r)}$  in time. This brings to the fore the pervasive issue of time scales in dynamical many-body systems, our specific question being the relative degree of localization ascribable to  $K_{SS'}^{(p)}$  and  $K_{SS'}^{(r)}$ . If the laser radiation is localized as a pulse, it can still extend over a period of time much longer than the extent of  $K_{SS'}^{(p)}(t)$ . In other words, the delta function representing  $K_{SS'}^{(p)}(t)$  may be quite different from the delta function representing  $K_{SS'}^{(r)}(t)$ , and the former may in fact not really be considered as localized with respect to the latter. With these provisos in mind, we can proceed to write the Markovian version of Eqs. (14), (17), (19) and (20):

$$\underline{K}(t) = (\underline{\Omega} + \underline{f}) \delta(t) , \quad (23)$$

$$\dot{\underline{P}}(t) = \underline{W} \cdot \underline{P}(t) , \quad (24)$$

$$\underline{J}(s) \cdot \underline{P}(s) = \underline{U} , \quad (25)$$

where

$$\underline{f} = \underline{k} \tau , \quad (26)$$

$$\underline{W} = \underline{\Omega} + \underline{f} , \quad (27)$$

$$\underline{J}(s) = (\underline{\Omega} + \underline{f} - s\underline{\Delta}) , \quad (28)$$

$$\underline{U} = - \underline{P}(0) , \quad (29)$$

and  $\tau$  represents the actual extent of the laser signal or a variable time parameter. Equation (24) is the well-known Pauli master equation (PME).

### III. Results of the Exact, Isomnesic and Markovian Approximations

The formalism of Sections I and II was applied to the system  $H(H_2O)/KCl(001)$ , the primary reasons for choosing which are the shallow adsorption potential well and the small number of bound states (7) it generates.<sup>17</sup> In Fig. 2 the results of the IA and the exact GME are compared for the system of references 11 and 15. The two are identical for  $H(H_2O)/KCl$ , and are not shown. While this is a short-time-scale comparison, the closeness

of the two leads us to expect the long-time IA results to be reliable. The Markovian approximation, on the other hand, cannot provide an unambiguous comparison because of large variations in the results for different values of  $\tau$ , shown in Fig. 3. All subsequent figures, therefore present our IA results.

Figure 4 is a composite display of the function  $E(t)$  obtained from results of Eq. (28) for six ranges of time differing by factors of 10 and for a laser intensity  $I = 1 \text{ W/cm}^2$ . It is seen that in going from a pico-second time range to a micro-second time range, the behaviour of the system changes dramatically. The early behaviour is dominated by fast energy transfer to phonon modes, completely overwhelming the slow laser pumping. The monotonic decay is representative of relaxation phenomena. For the 100 ps range, a false steady state seems to have been reached, but is in fact only an artifact of the large difference between  $A_r$  and  $A_p$ . The adbond energy starts to increase in the ns range, again suggesting a possible steady state. However, this is a transitional period during which the relative importance of the phonon terms and the laser terms starts to change. The latter, in fact, originate from a probability profile of the form  $(Ae^{\alpha t} \cos \beta t - Be^{\alpha t} \sin \beta t)$ , where  $\alpha + i\beta$  are complex roots of

$$\det \underline{M}(s) = 0, \quad (30)$$

and  $A + iB$  are complex coefficients obtained from  $\underline{M}(s)$  and  $\underline{I}(s)$ .

The phonon terms come from real roots and are therefore nonoscillatory. The magnitudes of the exponents  $\alpha$  in the laser part of the solution are much smaller than corresponding exponents in the phonon term. This is responsible for the radically different nature of  $E(t)$  in the different time regimes. Thus, the seemingly monotonic rise in  $E(t)$  apparent in the 10 ns regime is seen to be only the early segment of a cycle which

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

~~Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627~~

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Captain Lee Myers  
AFOSR/NC  
Bolling AFB  
Washington, D.C. 20332

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. G. D. Stein  
Mechanical Engineering Department  
Northwestern University  
Evanston, Illinois 60201

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G. H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. P. Hansma  
Physics Department  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
California Institute of Technology  
Division of Chemistry  
Pasadena, California 91125

Dr. W. Goddard  
California Institute of Technology  
Division of Chemistry  
Pasadena, California 91125

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6132  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Richard Colton  
Code 6112  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 5570  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
California Institute of Technology  
Division of Chemistry and Chemical  
Engineering  
Pasadena, California 91125

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
Southampton University  
Southampton SO9 5NH  
Hampshire, England

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

Dr. Richard Smardzewski  
Code 6130  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217



ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Surface Chemistry Division (6170)  
455 Overlook Avenue, S.W.  
Washington, D.C. 20375

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. P. Lund  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

DL/413/83/01  
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1




Fig. 5





Fig. 4



# EXACT vs. ISOMNESIC




Fig. 2




Fig. 1

Figure Captions

- Fig. 1. Memory kernels  $K_{SS'}^{(p)}(t)$  for  $(S, S') = (0, 3)$ . Solid line:  $3 \rightarrow 0$ ; dotted line:  $0 \rightarrow 3$ .
- Fig. 2. Probability profiles for system of references 11 and 15 obtained by solving the exact GME (solid line) and the isommesic GME (dashed line).
- Fig. 3. Average adbond energy  $E(t)$  obtained within the Markovian approximation for a range of values of the parameter  $\tau$ .
- Fig. 4.  $E(t)$  obtained by using the isommesic approximation for time scales ranging from picoseconds to microseconds, for laser intensity  $I = 1 \text{ W/cm}^2$ .
- Fig. 5. Same as Fig. 4 for  $I = 25 \text{ W/cm}^2$ .



16. W. H. Louisell, "Quantum Statistical Properties of Radiation"  
(John Wiley & Sons, New York 1973), Chapter 6.
17. H. Frank, H. Hoinkes and H. Wilsch, Surf. Sci. 63, 121 (1977).

## References

1. T. F. George, A. C. Beri, K. S. Lam and J. Lin, in Laser Applications, edited by J. F. Ready and R. K. Erf (Academic Press, New York, 1984), pp. 69-127.
2. T. F. George, J. Lin, A. C. Beri and W. C. Murphy, Prog. Surf. Sci. 16, 139 (1984).
3. T. J. Chuang, Surf. Sci. Rep. 3, 1 (1983).
4. J. Lin and T. F. George, J. Chem. Phys. 72, 2554 (1980);  
J. Lin and T. F. George, Phys. Rev. B 24, 64 (1981).
5. J. Lin, A. C. Beri, M. Hutchinson, W. C. Murphy and T. F. George, Phys. Lett. 79A, 233 (1980).
6. A. C. Beri and T. F. George, J. Chem. Phys. 78, 4288 (1983).
7. H. J. Kreuzer and D. N. Lowy, Chem. Phys. Lett. 78, 50 (1981); D. Lucas and G. E. Ewing, Chem. Phys. 58, 385 (1981); Z. W. Gortel, H. J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 27, 5066 (1983); 28, 2119 (1983).
8. C. Jedrzejek, K. F. Freed, S. Efrima and H. Metiu, Surf. Sci. 109, 191 (1981).
9. T. J. Chuang and H. Seki, Phys. Rev. Lett. 49, 382; T. J. Chuang, J. Chem. Phys. 76, 3828 (1982).
10. W. C. Murphy and T. F. George, Surf. Sci. 102, L46 (1981).
11. A. C. Beri and T. F. George, J. Chem. Phys., in press.
12. R. W. Zwanzig, in Lectures in Theoretical Physics, Vol. III, edited by W. E. Brittin, B. W. Downs and J. Downs (Interscience Publishers, New York 1961), p. 106ff.
13. F. Haake, in Springer Tracts in Modern Physics, edited by G. Höhler (Springer-Verlag, Berlin, 1973), p. 98ff.
14. C. Jedrzejek, K. F. Freed, E. Hood and H. Metiu, J. Chem. Phys. 79, 2436 (1983).
15. A. C. Beri and T. F. George, unpublished.

result, the sharp transitions assumed here will be broadened by a detuning-type effect, and estimates of the desorption rate presented above will have to be modified. However, the selectivity represented by the dominant  $T_1$  energy-transfer process will be retained.

For a complete study of desorption, transitions to the continuum must be included, and at the outset it is difficult to predict which levels will be most important. Work in this direction is in progress, and represents a departure from the first-principles approach in the inclusion of the bound-to-continuum transitions. This, as well as a complete first-principles treatment of vibrational LSSP including desorption, will be presented in the near future.

#### Acknowledgments

This work was supported in part by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Grant AFOSR-82-0046. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. TFG acknowledges the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Award (1974-86).

becomes evident in the 100 ns results and beyond. The frequency of this oscillation is seen to be  $k_0^{\frac{1}{2}}$ , where  $k_0$  is the amplitude of the laser kernel  $k_{SS}$ , and is essentially the Rabi frequency for the pair of levels resonant with the laser.

While the frequency of the long-term oscillations of  $P(t)$  and  $E(t)$  depends only on the laser term, being proportional to the square root of the intensity, the rate of onset of this behaviour depends on  $\alpha$  which varies linearly as  $k_0$  and inversely as the amplitude  $\Omega_0$  of the phonon kernel. Thus, increasing the laser power from 1 W/cm<sup>2</sup> to 25 W/cm<sup>2</sup> advances the onset time for the oscillatory state while increasing the frequency by a factor of 5, as displayed in Fig. 5.

Physically, the results of Figs. 4 and 5 can be interpreted in terms of desorption rates. Thus, the time required for  $|A|e^{\alpha t}$  and  $|B|e^{\alpha t}$  to become large enough so that the amplitude of  $E(t)$  in the oscillatory regime is substantially larger than the  $(t = 0)$  equilibrium Boltzmann value, is a reasonable measure of the desorption time to within an order of magnitude. This time is estimated by  $\tau_D = \alpha^{-1}$ . For our case,

$$\tau_D^{-1} = \alpha \approx .5 \frac{k_0}{\Omega_0} \approx (10^4 w) s^{-1}, \quad (31)$$

where  $w$  is laser power in W/cm<sup>2</sup>. Thus, for a 1 W/cm<sup>2</sup> laser, a hydrogen atom will stay adsorbed on the KCl(H<sub>2</sub>O) surface for  $\sim 10^{-4}$  s on the average before desorbing, or at least becoming very highly excited. The data mentioned here is for  $T = 150$  K.

It must be pointed out that a number of mechanisms whereby the desorption cross section could be modified are not included. Spontaneous decay, phase relaxation, phonon-phonon interaction and electron-hole pair creation are some of the ones being considered in extensions of this work. As a

DL/413/83/01  
056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

**END**

**FILMED**

**6-85**

**DTIC**